

NARRATOR: ALPROLIX® [Coagulation Factor IX (Recombinant), Fc Fusion Protein].

CASEY: My name's Casey, I'm from Ohio. I have hemophilia B. I have a beautiful wife named Mary and two beautiful daughters, Millie and Lucy, and I am an ALPROLIX peer, and I am also a pinsetter mechanic at a bowling alley. My career started when I was about 14 years old. My uncle and father both owned a family bowling alley business. I never got to see my dad a lot when I was younger. He gave me my treatments at 3-4:00 in the morning, because he worked the late shift. So when I was old enough to start working at the bowling alley, I started working out there so I could be around him more. So early in my career working in the bowling alley, I was always concerned about getting hurt by maybe a spring falling off or maybe heavy object hitting me, somebody hitting me with a bowling ball, but I loved doing it, I loved fixing machines. After talking to one of my nurses at the hemophilia center about ALPROLIX and its long half life, we decided to look over all the possible side effects and benefits of it, and then from there, we decided to start taking it. My doctor told me that allergic reactions may occur with ALPROLIX. My hemophilia center started me on Prophy doses every 10 days, and then they adjusted my dose based upon my response. Prophy dosing with ALPROLIX helps keep me protected from bleeds while I'm working on the job. The MyALPROLIX™ support team has been very helpful. They called my insurance, and then they got right back to me and told me that I was covered for four months. They were very helpful at getting me my factor product, answering any questions of mine, I would describe them as kind, helpful, and very, very supportive. I am very, very happy with the results. This is my experience, and everyone is different.

NARRATOR: Indications: ALPROLIX® [Coagulation Factor IX (Recombinant), Fc Fusion Protein] is a recombinant DNA derived coagulation factor IX concentrate indicated in adults and children with hemophilia B for on-demand treatment and control of bleeding episodes, perioperative management of bleeding, routine prophylaxis to reduce the frequency of bleeding episodes. Limitation of use: ALPROLIX is not indicated for induction of immune tolerance in patients with hemophilia B.

Important Safety Information: Do not use ALPROLIX if you are allergic to ALPROLIX or any of the other ingredients in ALPROLIX.

Tell your healthcare provider if you have or have had any medical problems, take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines, have any allergies and all your medical conditions, including if you are pregnant or planning to become pregnant, are breastfeeding, or have been told you have inhibitors (antibodies) to Factor IX.

Common side effects of ALPROLIX include headache, abnormal sensation in the mouth, and pain in your side with blood in your urine, which may be a sign of clot formation in the urinary collecting system. These are not all of the possible side effects of ALPROLIX. Talk to your

healthcare provider right away about any side effect that bothers you or does not go away, and if bleeding is not controlled using ALPROLIX.

Allergic reactions may occur with ALPROLIX. Call your healthcare provider or get emergency treatment right away if you have any of the following symptoms: hives, chest tightness, wheezing, difficulty breathing, or swelling of the face. Your body can also make antibodies called "inhibitors" against ALPROLIX, which may stop ALPROLIX from working properly.

ALPROLIX may increase the risk of formation of abnormal blood clots in your body, especially if you have risk factors for developing blood clots. Call your healthcare provider or seek emergency care if you have symptoms of a possible abnormal blood clot, which may include: chest pain, difficulty breathing, unexpected swelling of an arm or leg with or without pain or tenderness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see Full Prescribing Information.

[00:05:00]